

- 60 **Superparamagnetic iron oxide-enhanced MRI at 3 T for accurate axillary staging in breast cancer**  
*K. Motomura, T. Izumi, S. Tateishi, Y. Tamaki, Y. Ito, T. Horinouchi and K. Nakanishi*  
 Imaging replaces sentinel node biopsy
- 70 **Radioactive seed localization in breast cancer treatment**  
*N. N. Y. Janssen, J. Nijkamp, T. Alderliesten, C. E. Loo, E. J. T. Rutgers, J.-J. Sonke and M. T. F. D. Vrancken Peeters*  
 Facilitates logistics
- 81 **Pooled long-term outcomes from two randomized trials of axillary node sampling with axillary radiotherapy versus axillary node clearance in patients with operable node-positive breast cancer**  
*A. U. Bing, G. R. Kerr, W. Jack, U. Chetty, L. J. Williams, A. Rodger and J. M. Dixon*  
 No difference in outcome
- 88 **National proficiency-gain curves for minimally invasive gastrointestinal cancer surgery**  
*H. Mackenzie, S. R. Markar, A. Askari, M. Ni, O. Faiz and G. B. Hanna*  
 Numbers needed to learn
- 97 **Impact of number of [<sup>18</sup>F]fluorodeoxyglucose-PET-positive lymph nodes on survival of patients receiving neoadjuvant chemotherapy and surgery for oesophageal cancer**  
*H. Miyata, M. Yamasaki, T. Makino, M. Tatsumi, Y. Miyazaki, T. Takahashi, Y. Kurokawa, S. Takiguchi, M. Mori and Y. Doki*  
 Dismal prognosis if PET-positive nodes exist after treatment
- 105 **Risk adjustment models for short-term outcomes after surgical resection for oesophagogastric cancer**  
*C. Fischer, H. Lingsma, R. Hardwick, D. A. Cromwell, E. Steyerberg and O. Groene*  
 Needs to be tested outside England and Wales
- 117 **Patterns of recurrence in early-stage oesophageal cancer after chemoradiotherapy and surgery compared with surgery alone**  
*W. B. Robb, M. Messenger, L. Dahan, F. Mornex, E. Maillard, X. B. D'Journo, J.-P. Triboulet, L. Bedenne, J.-F. Seitz and C. Mariette, on behalf of the Fédération Francophone de Cancérologie Digestive, Société Française de Radiothérapie Oncologique, Union des Centres de Lutte Contre le Cancer, Groupe Coopérateur Multidisciplinaire en Oncologie and the French EsoGAstric Tumour working group – Fédération de Recherche En Chirurgie*  
 Multimodal treatment reduces locoregional but not distant metastatic recurrence
- 126 **Radiofrequency ablation as an alternative to hepatic resection for single small hepatocellular carcinomas**  
*G.-A. Kim, J. H. Shim, M.-J. Kim, S. Y. Kim, H. J. Won, Y. M. Shin, P. N. Kim, K.-H. Kim, S.-G. Lee and H. C. Lee*  
 Good alternative in small hepatocellular carcinoma
- 136 **Volume–outcome relationship in pancreatic surgery**  
*G. Alsfasser, H. Leicht, C. Günster, B. M. Rau, G. Schillinger and E. Klar*  
 Better outcomes after 1 year in high-volume hospitals